

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Transcriptional study of cerato-platanin encoding genes in homokaryotic and heterokaryotic isolates of the forest pathogen *Heterobasidion irregulare*

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1551810> since 2016-06-24T17:25:40Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution:

Questa è la versione dell'autore dell'opera:

[Baccelli I., Gonthier P., Bernardi R., 2015. Journal of Plant Pathology, 97, S40]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://sipav.org/main/jpp/index.php/jpp>]

**TRANSCRIPTIONAL STUDY OF CERATO-PLATANIN
ENCODING GENES IN HOMOKARYOTIC AND HETEROKARYOTIC
ISOLATES OF THE FOREST PATHOGEN**

HETEROBASIDION IRREGULARE. I. Baccelli¹, P. Gonthier²,

R. Bernardi³. ¹Department of Agrifood Production and Environmental Sciences (DISPAA), University of Firenze, Piazzale delle Cascine 28, I-50144 Firenze, Italy. ²Department of Agricultural, Forest and Food Sciences (DISAFA), University of Torino, Largo Paolo Braccini 2, I-10095 Grugliasco (TO), Italy. ³Department of Agriculture, Food and Environment (DISAAA-a), University of Pisa, Via del Borghetto 80, I-56124 Pisa, Italy. E-mail: ivan.baccelli@inwind.it

The forest pathogen *Heterobasidion irregulare* (Basidiomycota) was recently sequenced and three cerato-platanin encoding genes were found in its genome (*HiCPs*). Cerato-Platanin Proteins (CPPs) are a family of proteins well known for their resistance-inducing ability when applied on plants. What is still poorly understood is their role in fungal life. These proteins seem to play both a role in the fungal cell wall and a role in the fungus-plant interaction, but most data available to date on CPPs derive from studies performed on Ascomycetes. In the present study, we investigated the expression of *HiCPs* in three homokaryotic isolates and two heterokaryotic isolates of the plant pathogen *H. irregulare*. Homokaryotic and heterokaryotic mycelia not only play different roles in the infectious process but also differ in their biology: clamp connections are formed in the heterokaryotic mycelia. Transcription of *HiCPs* was analysed both at the edge and at the centre of the fungal colony and compared between homokaryon and heterokaryon. Results showed *HiCP1* to be the gene with the highest transcript abundance among *HiCPs*. *HiCP1* did not show any preferential expression in different sections of the fungal colony, while *HiCP2* was significantly more expressed at the colony centre, thus suggesting a link with the production of conidia. The level of expression of *HiCPs* in heterokaryons was generally comparable to that of one or both the parental homokaryons, irrespective of the colony section, thus demonstrating that *HiCPs* are not transcriptionally influenced by the heterokaryotic stage.